

POLYNOMIAL PRESERVING GRADIENT RECOVERY AND A *POSTERIORI* ESTIMATE FOR BILINEAR ELEMENT ON IRREGULAR QUADRILATERALS

ZHIMIN ZHANG

Abstract. A polynomial preserving gradient recovery method is proposed and analyzed for bilinear element under quadrilateral meshes. It has been proven that the recovered gradient converges at a rate $O(h^{1+\rho})$ for $\rho = \min(\alpha, 1)$, when the mesh is distorted $O(h^{1+\alpha})$ ($\alpha > 0$) from a regular one. Consequently, the *a posteriori* error estimator based on the recovered gradient is asymptotically exact.

Key Words. Finite element method, quadrilateral mesh, gradient recovery, superconvergence, *a posteriori* error estimate.

1. Introduction

A posteriori error estimation is an active research area and many methods have been developed. Roughly speaking, there are residual type error estimators and recovery type estimators. For the literature, readers are referred to recent books by Ainsworth-Oden [2] and by Babuška-Strouboulis [4], a conference proceeding [16], a survey article by Bank [5], an earlier book by Verfürth [23], and references therein.

While residual type estimators have been analyzed extensively, there is only limited theoretical research on recovery type error estimators (see, e.g., [2, Chapter 4], [6, 7, 9, 10, 15, 22, 28, 29]). Yet, recovery type error estimators are widely used in engineering applications and their practical effectiveness has been recognized by more and more researchers. Currently, *ZZ* patch recovery is used in commercial codes, such as ANSYS, MCS/NASTRAN-Marc, Pro/MECHANICA (a product of Parametric Technology), and IDEAS (a product of SDRC, part of EDS), for the purpose of smoothing and adaptive re-meshing. It is also used in NASA's COMET-AR (COMputational MEchanics Testbed With Adaptive Refinement). In a computer based investigation [4] by Babuška et al., it was found that among all error estimators tested (including the equilibrated residual error estimator, the *ZZ* patch recovery error estimator, and many others), the *ZZ* patch recovery error estimator based on the discrete least-squares fitting is the most robust.

2000 *Mathematics Subject Classification.* 65N30, 65N15, 41A10, 41A25, 41A27, 41A63.

This research was partially supported by the National Science Foundation grants DMS-0074301 and DMS-0311807.

It is worth pointing out that the recovery type error estimator was originally based on finite element superconvergence theory, in hopes that a recovered gradient was superconvergent and hence could be used as a substitute of the exact gradient to measure the error. The reader is referred to [4, 11, 16, 18, 25, 34] for literature regarding superconvergence theory. In order to prove superconvergence, it is necessary to impose some strong restrictions on mesh, which are usually not satisfied in practice. Nevertheless, it is found that in many practical situations, recovery type error estimators perform astonishingly well under meshes produced by the Delaunay triangulation. Mathematically, this fact has not yet been rigorously justified.

In a recent work, Bank-Xu [6, 7] introduced a recovery type error estimator based on global L_2 -projection with smoothing iteration of the multigrid method, and they established asymptotic exactness in the H^1 -norm for linear element under shape regular triangulation. However, the recovery operator is a global one.

On the other hand, Wang proposed a “semi-local” recovery [27] and proved its superconvergence under the quasi-uniform mesh assumption. The main feature of his method is to apply L_2 projection on a coarser mesh with size $\tau = Ch^\alpha$ with $\alpha \in (0, 1)$. Consequently, there is no upper bound for the number of elements in an element patch when mesh size $h \rightarrow 0$.

As for element-wise recovery operators, Schatz-Wahlbin et al. [15, 22] established a general framework which requests, for linear element, given a fixed $0 < \epsilon < 1$, that

$$m = C \left(\left(\frac{H}{h} \right)^2 h^\epsilon + \left(\frac{h}{H} \right)^\epsilon \ln \frac{H}{h} \right) < 1.$$

Here h is the size of element τ , $H \geq 2h$ is the size of the patch ω_τ (surrounding τ), where the recovery takes place, and C is an unknown constant which comes from the analysis. Let $H = Lh$. In order for $m < 1$, we need

$$C(L^2 h^\epsilon + L^{-\epsilon} \ln L) < 1.$$

Depending on C , this essentially asks for sufficiently large L and sufficiently small h , which implies many elements may be needed for the recovery operator. Nevertheless, in practice, many recovery operators work well with an H/h that is not large (usually 2).

Therefore a theoretical justification for recovery that involves only a few elements surrounding a node is necessary. In other word, it is desired to study the case when $H = 2h$. The situation is further complicated by quadrilateral meshes where mappings between the reference element and physical elements are not affine. We encounter some delicate theoretical issue in analysis. See [1, 3, 8, 13, 14, 19, 21, 30, 31, 37] for more details.

In this article, we propose and analyze a gradient recovery method which is different from the ZZ recovery [35]. We show that the *a posteriori* estimate based on this new recovery operator is asymptotically exact under distortion $O(h^{1+\alpha})$ when $\alpha > 0$ for quadrilateral meshes. Here $\alpha = \infty$ represents the uniform mesh and $\alpha = 0$ represents completely unstructured mesh. For counterpart results under triangular meshes, the reader is referred to [32, 33].

Due to non-affine mapping feature of the quadrilateral mesh, the analysis is much harder than the triangular mesh.

The main feature of this new recovery operator is:

- (1) It is completely local just like the ZZ patch recovery;
- (2) It is polynomial preserving under practical meshes, a property not shared by the ZZ;
- (3) It is superconvergent under minorly restricted mesh conditions;
- (4) It results in an asymptotically exact error estimator when the mesh is not overly distorted. The error bound is in the form of

$$(1.1) \quad \eta_h + O(h^{1+\rho}) \leq \|\nabla(u - u_h)\| \leq \eta_h + O(h^{1+\rho}),$$

rather than

$$\frac{1}{C}\eta_h + \text{higher order term} \leq \|\nabla(u - u_h)\| \leq C\eta_h + \text{higher order term}$$

in most error bounds in the literature. Here C is an unknown constant, which may be very large and hence makes the error bound not very meaningful in practice.

We comment that h^α can be reduced to $o(1)$ and still maintain the asymptotic exactness of the error estimator. If we give up the asymptotic exactness requirement and only ask for a reasonable error estimator, we may further reduce the condition to “a sufficiently small constant $\gamma > 0$ ”.

The main results of this paper include a super-close property (Theorem 3.3), a global superconvergent recovery result (Theorem 4.2), and a local superconvergent recovery result (Theorem 4.3). The error bound (1.1) is a consequence of Theorems 4.2 or 4.3. All these results need a mesh assumption, *Condition* (α), which is introduced in Section 2. Basically, we allow quadrilaterals to be asymptotically distorted by $O(h^{1+\alpha})$ ($\alpha > 0$) from parallelograms. Note that $\alpha = 0$ represents arbitrary meshes. Therefore, the mesh considered here is next to arbitrary (with a little structure). Indeed, when a very practical mesh refinement strategy, bisection (link edge-center of each opposite side of a quadrilateral) is applied, we have $\alpha = 1$ (see Lemma 2.1).

2. Geometry of the Quadrilateral

Let $\hat{K} = [-1, 1] \times [-1, 1]$ be the reference element with vertices \hat{Z}_i , and let K be a convex quadrilateral with vertices $Z_i^K(x_i^K, y_i^K)$, $i = 1, 2, 3, 4$. There exists a unique bilinear mapping F_K such that $F_K(\hat{K}) = K$, $F_K(\hat{Z}_i) = Z_i^K$ given by

$$x = \sum_{i=1}^4 x_i^K N_i, \quad y = \sum_{i=1}^4 y_i^K N_i,$$

where

$$\begin{aligned} N_1 &= \frac{1}{4}(1 - \xi)(1 - \eta), & N_2 &= \frac{1}{4}(1 + \xi)(1 - \eta), \\ N_3 &= \frac{1}{4}(1 + \xi)(1 + \eta), & N_4 &= \frac{1}{4}(1 - \xi)(1 + \eta). \end{aligned}$$

We can also express

$$x = a_0 + a_1\xi + a_2\eta + a_3\xi\eta, \quad y = b_0 + b_1\xi + b_2\eta + b_3\xi\eta;$$

where by suppressing the index “ K ”,

$$\begin{aligned} 4a_0 &= x_1 + x_2 + x_3 + x_4, & 4b_0 &= y_1 + y_2 + y_3 + y_4; \\ 4a_1 &= -x_1 + x_2 + x_3 - x_4, & 4b_1 &= -y_1 + y_2 + y_3 - y_4; \\ 4a_2 &= -x_1 - x_2 + x_3 + x_4, & 4b_2 &= -y_1 - y_2 + y_3 + y_4; \\ 4a_3 &= x_1 - x_2 + x_3 - x_4, & 4b_3 &= y_1 - y_2 + y_3 - y_4. \end{aligned}$$

To any function $v(x, y)$ defined on K , we associate $\hat{v}(\xi, \eta)$ by

$$\hat{v}(\xi, \eta) = v(x(\xi, \eta), y(\xi, \eta)), \quad \text{or} \quad \hat{v} = v \circ F_K.$$

The Jacobi matrix of the mapping F_K is

$$(DF_K)(\xi, \eta) = \begin{pmatrix} x_\xi & y_\xi \\ x_\eta & y_\eta \end{pmatrix} = \begin{pmatrix} a_1 + a_3\eta & b_1 + b_3\eta \\ a_2 + a_3\xi & b_2 + b_3\xi \end{pmatrix}.$$

Let $\nabla v = (\partial_x v, \partial_y v)^T$, it is straight forward to verify that

$$(2.1) \quad \hat{\nabla} \hat{v} = (\partial_\xi \hat{v}, \partial_\eta \hat{v})^T = DF_K \nabla v,$$

$$(2.2) \quad \partial_{\xi^r}^r \hat{v} = [(a_1 + a_3\eta)\partial_x + (b_1 + b_3\eta)\partial_y]^r v,$$

$$(2.3) \quad \begin{aligned} \partial_{\xi^r \eta}^{r+1} \hat{v} &= r[(a_1 + a_3\eta)\partial_x + (b_1 + b_3\eta)\partial_y]^{r-1} (a_3, b_3) \cdot \nabla v \\ &\quad + [(a_1 + a_3\eta)\partial_x + (b_1 + b_3\eta)\partial_y]^r [(a_2 + a_3\xi)\partial_x + (b_2 + b_3\xi)\partial_y] v, \end{aligned}$$

and $\partial_{\eta^r}^r \hat{v}$ and $\partial_{\xi \eta^r}^{r+1} \hat{v}$ can be expressed in a similar way. The determinant of the Jacobi matrix is

$$J_K = J_K(\xi, \eta) = J_0^K + J_1^K \xi + J_2^K \eta,$$

where

$$J_0^K = a_1 b_2 - b_1 a_2, \quad J_1^K = a_1 b_3 - b_1 a_3, \quad J_2^K = b_2 a_3 - a_2 b_3.$$

The inverse of the Jacobi matrix is

$$\begin{pmatrix} \xi_x & \eta_x \\ \xi_y & \eta_y \end{pmatrix} = (DF_K)^{-1} = \frac{1}{J_K} \begin{pmatrix} b_2 + b_3\xi & -b_1 - b_3\eta \\ -a_2 - a_3\xi & a_1 + a_3\eta \end{pmatrix}.$$

Note that $a_3 = b_3 = 0$ when K is a parallelogram in which case F_K is an affine mapping, and further $a_3 = b_3 = a_2 = b_1 = 0$ when K is a rectangle.

Starting from Z_1 , we express the four edges (with the midpoint P_i) as four vectors \mathbf{v}_i , $i = 1, 2, 3, 4$, pointing counter-clock-wisely (Figure 1). We denote the midpoints of Z_2Z_4 and Z_1Z_3 as O_1 and O_2 , respectively. For analysis purpose, it is convenient to identify 2-D vectors as 3-D vectors by adding the third component 0. We can verify that

$$\begin{aligned} P_4P_2 &= \frac{1}{2}(x_2 + x_3 - x_4 - x_1, y_2 + y_3 - y_4 - y_1, 0) = 2(a_1, b_1, 0), \\ P_1P_3 &= \frac{1}{2}(x_3 + x_4 - x_1 - x_2, y_3 + y_4 - y_1 - y_2, 0) = 2(a_2, b_2, 0), \\ O_1O_2 &= \frac{1}{2}(x_1 + x_3 - x_2 - x_4, y_1 + y_3 - y_2 - y_4, 0) = 2(a_3, b_3, 0). \end{aligned}$$




FIGURE 1. Geometry of a quadrilateral

Then

$$(2.4) \quad 2\sqrt{a_1^2 + b_1^2} = |P_4P_2|, \quad 2\sqrt{a_2^2 + b_2^2} = |P_1P_3|, \quad 2\sqrt{a_3^2 + b_3^2} = |O_1O_2|.$$

$$(2.5) \quad 4(a_1a_2 + b_1b_2) = P_4P_2 \cdot P_1P_3 = |P_4P_2||P_1P_3| \cos \alpha_K,$$

$$(2.6) \quad 4(a_1a_3 + b_1b_3) = P_4P_2 \cdot O_1O_2 = |P_4P_2||O_1O_2| \cos \beta_K,$$

$$(2.7) \quad 4(a_2a_3 + b_2b_3) = O_1O_2 \cdot P_1P_3 = |O_1O_2||P_1P_3| \cos \gamma_K,$$

where the meaning of angles α_K , β_K , and γ_K is obvious from the context.

$$(2.8) \quad J_0^K \mathbf{k} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & b_1 & 0 \\ a_2 & b_2 & 0 \end{vmatrix} = \frac{1}{4} P_4P_2 \times P_1P_3 = \frac{1}{4} |P_4P_2||P_1P_3| \sin \alpha_K,$$

$$(2.9) \quad J_1^K \mathbf{k} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & b_1 & 0 \\ a_3 & b_3 & 0 \end{vmatrix} = \frac{1}{4} P_4P_2 \times O_1O_2 = \frac{1}{4} |P_4P_2||O_1O_2| \sin \beta_K,$$

$$(2.10) \quad J_2^K \mathbf{k} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_2 & b_2 & 0 \\ a_3 & b_3 & 0 \end{vmatrix} = \frac{1}{4} P_1P_3 \times O_1O_2 = \frac{1}{4} |P_1P_3||O_1O_2| \sin \gamma_K.$$

We could also express

$$|J_1^K| = 2|(x_4 - x_3)(y_2 - y_1) - (x_2 - x_1)(y_4 - y_3)| = 2|\mathbf{v}_3 \times \mathbf{v}_1|, \quad |J_2^K| = 2|\mathbf{v}_4 \times \mathbf{v}_2|.$$

Let h_K be the longest edge length of K , we introduce the following condition:

Definition 1. A convex quadrilateral K is said to satisfy the *diagonal condition* if

$$(2.11) \quad d_K = |O_1O_2| = O(h_K^{1+\alpha}), \quad \alpha \geq 0.$$

Note that K is a parallelogram if and only if $d_K = 0$. Therefore, the distance between the two diagonal mid-points O_1 and O_2 is a convenient measure for the deviation of a quadrilateral from a parallelogram. The two extremal cases $\alpha \rightarrow \infty$ and $\alpha \rightarrow 0$ represent parallelogram and completely unstructured quadrilateral, respectively. Anything in between will pose some restriction, especially $\alpha = 1$ is the well-known 2-strongly regular partition, see, e.g., [13, 37].

The diagonal condition was previously used by Chen [12] for triangular meshes, where two adjacent triangles form a quadrilateral that satisfies the condition.

The following lemma states a known fact regarding the 2-strongly regular partition ($\alpha = 1$). Although this fact is widely used, we have not seen a formal proof of it in the literature. An elementary proof is therefore provided in the Appendix.

Lemma 2.1. Let $o_1 o_2$ be the distance between two diagonal mid-points of any of four refined quadrilaterals through the bi-section of K . Then

$$|o_1 o_2| = \frac{1}{4} |O_1 O_2|.$$

Recall that the bi-section reduces the length of longest edge by half, which is $h_K/2$. Therefore, the diagonal condition (2.11) is satisfied with $\alpha = 1$.

To measure this deviation, Rannachar and Turek [21] used the quantity

$$\sigma_K = \max(|\pi - \theta_1|, |\pi - \theta_2|),$$

where θ_1 and θ_2 are the angles between the outward normals of two opposite sides of K .

Definition 2. A convex quadrilateral K is said to satisfy the *angle condition* if

$$(2.12) \quad \sigma_K = O(h_K^\alpha), \quad \alpha \geq 0.$$

Lemma 2.2. The diagonal condition (2.11) and the angle condition (2.12) are equivalent in the sense

$$d_K = O(h_K^{1+\alpha}) \iff \sigma_K = O(h_K^\alpha), \quad \alpha \geq 0.$$

A special case of this lemma has been proved in [19, Theorem 4.13] under some complicated mesh restrictions. Here we provide a direct and much simpler proof in the Appendix without any mesh assumption.

Definition 3. A partition \mathcal{T}_h is said to satisfy *Condition* (α) if there exist $\alpha > 0$ such that

- i) Any $K \in \mathcal{T}_h$ satisfies the diagonal condition (2.11).
- ii) Any two K_1, K_2 in \mathcal{T}_h that share a common edge satisfy a *neighboring condition*: For $j = 1, 2$,

$$(2.13) \quad a_j^{K_1} = a_j^{K_2} (1 + O(h_{K_1}^\alpha + h_{K_2}^\alpha)), \quad b_j^{K_1} = b_j^{K_2} (1 + O(h_{K_1}^\alpha + h_{K_2}^\alpha)).$$

To assure optimal order error estimates in the H^1 -norm for the bilinear isoparametric interpolation on a convex quadrilateral K , namely, the

estimate

$$(2.14) \quad \|u - u_I\|_{0,K} + h|u - u_I|_{1,K} \leq Ch_K^2 |u|_{2,K},$$

we need a degeneration condition, which was introduced by Acosta and Durán [1].

Definition 4. A convex quadrilateral K is said to satisfy the *Regular decomposition property* with constants $N \in \mathbb{R}$ and $0 < \Psi < \pi$, or shortly $RDP(N, \Psi)$, if we can divide K into two triangles along one of its diagonals, which will always be called d_1 , in such a way that $|d_1|/|d_2| \leq N$ and both triangles satisfy the *maximum angle condition* with parameter Ψ (i.e., all angles are bounded by Ψ).

Remark. This is a weaker condition than many other similar degenerate conditions, cf. e.g., [13, 14, 31, 37]. It was proved in [1] that $RDP(N, \Psi)$ is a sufficient condition for (2.14) to be hold, and the authors conjectured that it is also a necessary condition. Recently, Ming-Shi confirmed this conjecture by a simple counter-example [19].

We denote $X = X(\xi, \eta) = X_0 + X_1$ where

$$X_0 = \begin{pmatrix} b_2 & -b_1 \\ -a_2 & a_1 \end{pmatrix}, \quad X_1 = X_1(\xi, \eta) = \begin{pmatrix} b_3 \\ -a_3 \end{pmatrix} (\xi, -\eta).$$

Lemma 2.3. Let a convex quadrilateral K satisfy the diagonal condition. Then

$$\|X_0 X^{-1}\|_2 = 1 + O(h_K^\alpha), \quad \|X_1 X^{-1}\|_2 = \|I - X_0 X^{-1}\|_2 = O(h_K^\alpha).$$

Proof: It is straightforward to verify that

$$\begin{aligned} X_0 X^{-1} &= \begin{pmatrix} b_2 & -b_1 \\ -a_2 & a_1 \end{pmatrix} \frac{1}{J_K} \left[\begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} + \begin{pmatrix} \eta \\ \xi \end{pmatrix} (a_3, b_3) \right] \\ &= \frac{J_0^K}{J_K} I + \frac{1}{J_K} \begin{pmatrix} b_2 & -b_1 \\ -a_2 & a_1 \end{pmatrix} \begin{pmatrix} \eta \\ \xi \end{pmatrix} (a_3, b_3) \end{aligned}$$

where I is a 2-by-2 identity matrix; and

$$X_1 X^{-1} = I - X_0 X^{-1} = \left(\frac{J_1^K}{J_K} \xi + \frac{J_2^K}{J_K} \eta \right) I - \frac{1}{J_K} \begin{pmatrix} b_2 & -b_1 \\ -a_2 & a_1 \end{pmatrix} \begin{pmatrix} \eta \\ \xi \end{pmatrix} (a_3, b_3).$$

By the definition of J_K and geometric relations of (2.4), (2.8)–(2.10), we see that

$$\frac{J_0^K}{J_K} = 1 + O(h_K^\alpha), \quad \frac{J_1^K}{J_K} = O(h_K^\alpha), \quad \frac{J_2^K}{J_K} = O(h_K^\alpha),$$

by the diagonal condition (2.11). The desired conclusion follows. \square

3. Superconvergence Analysis

We consider the variational problem: Find $u \in H^1(\Omega)$ such that

$$(3.1) \quad a(u, v) = (\nabla u, A \nabla v) + (\mathbf{b} \cdot \nabla u, v) + (cu, v) = (f, v), \quad \forall v \in H^1(\Omega),$$

where A is a 2-by-2 symmetric positive definite matrix and Ω is a polygonal domain which allows a quadrilateral partition \mathcal{T}_h with $h = \max_{K \in \mathcal{T}_h} h_K$. We assume that all functions are sufficiently smooth, in particular,

$$(3.2) \quad \|A - A_0\|_{0,\infty,K} = O(h_K^\alpha), \quad \|\mathbf{b} - \mathbf{b}_0\|_{0,\infty,K} = O(h_K^\alpha),$$

where A_0 and \mathbf{b}_0 are piece-wisely constant functions that on each $K \in \mathcal{T}_h$,

$$A_0|_K = \frac{1}{|K|} \int_K A(x, y) dx dy, \quad \mathbf{b}_0|_K = \frac{1}{|K|} \int_K \mathbf{b}(x, y) dx dy.$$

We also assume that $a(\cdot, \cdot)$ satisfies the inf-sup condition to insure that (3.1) has a unique solution. Using

$$\nabla v = \frac{1}{J_K} X \hat{\nabla} \hat{v},$$

we write

$$\begin{aligned} (\nabla w, A \nabla v)_K &= \int_K (\nabla w)^T A \nabla v dx dy = \int_{\hat{K}} \frac{1}{J_K} (X \hat{\nabla} \hat{w})^T \hat{A} (X \hat{\nabla} \hat{v}) d\xi d\eta, \\ (\mathbf{b} \cdot \nabla w, v)_K &= \int_K v \mathbf{b} \cdot \nabla w dx dy = \int_{\hat{K}} \hat{v} \hat{\mathbf{b}} \cdot X \hat{\nabla} \hat{w} d\xi d\eta; \end{aligned}$$

and define

$$(3.3) \quad \begin{aligned} (\nabla w, A \nabla v)_K^* &= \frac{1}{J_0^K} \int_{\hat{K}} (X_0 \hat{\nabla} \hat{w})^T A_0 (X_0 \hat{\nabla} \hat{v}) d\xi d\eta \\ &= \int_{\hat{K}} (\hat{\nabla} \hat{w})^T B^K \hat{\nabla} \hat{v} d\xi d\eta, \end{aligned}$$

$$(3.4) \quad (\mathbf{b} \cdot \nabla w, v)_K^* = \mathbf{b}_0 \cdot X_0 \int_{\hat{K}} \hat{v} \hat{\nabla} \hat{w} d\xi d\eta,$$

where

$$B^K = \frac{1}{J_0^K} (X_0^K)^T A_0^K X_0^K.$$

We introduce the following lemma, which can be verified by straightforward calculation.

Lemma 3.1. Under the condition (2.13) and (3.2), we have

$$J_0^{K_1} = J_0^{K_2} (1 + O(h_{K_1}^\alpha + h_{K_2}^\alpha)), \quad \|B^{K_1} - B^{K_2}\| = O(h_{K_1}^\alpha + h_{K_2}^\alpha).$$

Theorem 3.1. Let the assumption (3.2) be satisfied, and let K satisfy the diagonal condition. Then there exists a constant C independent of u and K , such that

$$(3.5) \quad |(\nabla w, A \nabla v)_K - (\nabla w, A \nabla v)_K^*| \leq C h_K^\alpha \|\nabla w\|_{0,K} \|\nabla v\|_{0,K},$$

$$(3.6) \quad |(\mathbf{b} \cdot \nabla w, v)_K - (\mathbf{b} \cdot \nabla w, v)_K^*| \leq C h_K^\alpha \|\nabla w\|_{0,K} \|v\|_{0,K}.$$

Proof: We decompose

$$(3.7) \quad (\nabla w, A\nabla v)_K - (\nabla w, A\nabla v)_K^* = (\nabla w, (A - A_0)\nabla v)_K \\ + \int_{\hat{K}} \frac{1}{J_K} [(X\hat{\nabla}\hat{w})^T A_0(X\hat{\nabla}\hat{v}) - (X_0\hat{\nabla}\hat{w})^T A_0(X_0\hat{\nabla}\hat{v})] d\xi d\eta \\ + \int_{\hat{K}} \left(\frac{1}{J_K} - \frac{1}{J_0^K}\right) (X_0\hat{\nabla}\hat{w})^T A_0(X_0\hat{\nabla}\hat{v}) d\xi d\eta.$$

By (3.2)

$$(3.8) \quad |(\nabla w, (A - A_0)\nabla v)_K| \leq Ch_K^\alpha \|\nabla w\|_{0,K} \|\nabla v\|_{0,K}.$$

Using $X = X_0 + X_1$, we express

$$\int_{\hat{K}} \frac{1}{J_K} [(X\hat{\nabla}\hat{w})^T A_0(X\hat{\nabla}\hat{v}) - (X_0\hat{\nabla}\hat{w})^T A_0(X_0\hat{\nabla}\hat{v})] d\xi d\eta \\ = \int_{\hat{K}} \frac{1}{J_K} [(X_0\hat{\nabla}\hat{w})^T A_0(X_1\hat{\nabla}\hat{v}) + (X_1\hat{\nabla}\hat{w})^T A_0(X_0\hat{\nabla}\hat{v}) \\ + (X_1\hat{\nabla}\hat{w})^T A_0(X_1\hat{\nabla}\hat{v})] d\xi d\eta.$$

The first term can be estimated as

$$\left| \int_{\hat{K}} \frac{1}{J_K} (X_0\hat{\nabla}\hat{w})^T A_0(X_1\hat{\nabla}\hat{v}) d\xi d\eta \right| \\ = \left| \int_{\hat{K}} \left(\frac{1}{J_K} X\hat{\nabla}\hat{w}\right)^T X^{-T} X_0^T A_0 X_1 X^{-1} \left(\frac{1}{J_K} X\hat{\nabla}\hat{v}\right) J_K d\xi d\eta \right| \\ = \left| \int_K (\nabla w)^T (X_0 X^{-1})^T A_0 X_1 X^{-1} \nabla v dx dy \right| \\ \leq Ch_K^\alpha \|\nabla w\|_{0,K} \|\nabla v\|_{0,K}.$$

Note that $(X_0 X^{-1})^T A_0 X_1 X^{-1} = O(h_K^\alpha)$ by Lemma 2.3. The other two terms can be estimated similarly. Then we derive

$$(3.9) \quad \left| \int_{\hat{K}} \frac{1}{J_K} [(X\hat{\nabla}\hat{w})^T A_0(X\hat{\nabla}\hat{v}) - (X_0\hat{\nabla}\hat{w})^T A_0(X_0\hat{\nabla}\hat{v})] d\xi d\eta \right| \\ \leq Ch_K^\alpha \|\nabla w\|_{0,K} \|\nabla v\|_{0,K}.$$

Next

$$(3.10) \quad \left| \int_{\hat{K}} \left(\frac{1}{J_K} - \frac{1}{J_0^K}\right) (X_0\hat{\nabla}\hat{w})^T A_0(X_0\hat{\nabla}\hat{v}) d\xi d\eta \right| \\ = \left| \int_{\hat{K}} \left(1 - \frac{J_K}{J_0^K}\right) \left(\frac{1}{J_K} X\hat{\nabla}\hat{w}\right)^T X^{-T} X_0^T A_0 X_0 X^{-1} \left(\frac{1}{J_K} X\hat{\nabla}\hat{v}\right) J_K d\xi d\eta \right| \\ = \left| \int_K \left(-\frac{J_1^K}{J_0^K} \xi(x, y) - \frac{J_2^K}{J_0^K} \eta(x, y)\right) (\nabla w)^T (X_0 X^{-1})^T A_0 X_0 X^{-1} \nabla v dx dy \right| \\ \leq Ch_K^\alpha \|\nabla w\|_{0,K} \|\nabla v\|_{0,K}.$$

Note that by Lemma 2.3,

$$\frac{J_1^K}{J_0^K} \xi(x, y) + \frac{J_2^K}{J_0^K} \eta(x, y) = O(h_K^\alpha), \quad (X_0 X^{-1})^T A_0 X_0 X^{-1} = 1 + O(h_K^\alpha).$$

We then obtain (3.5) by applying (3.8)-(3.10) to the right hand side of (3.7).

Now we write the convection term as following:

$$(3.11) \quad \begin{aligned} & (\mathbf{b} \cdot \nabla w, v)_K - (\mathbf{b} \cdot \nabla w, v)_K^* \\ &= \int_{\hat{K}} \hat{v} \hat{\mathbf{b}} \cdot (X - X_0) \hat{\nabla} \hat{w} d\xi d\eta + \int_{\hat{K}} \hat{v} (\hat{\mathbf{b}} - \mathbf{b}_0) \cdot X_0 \hat{\nabla} \hat{w} d\xi d\eta. \end{aligned}$$

We estimate the two terms separately.

$$(3.12) \quad \begin{aligned} & \left| \int_{\hat{K}} \hat{v} \hat{\mathbf{b}} \cdot (X - X_0) \hat{\nabla} \hat{w} d\xi d\eta \right| \\ &= \left| \int_{\hat{K}} \hat{v} \hat{\mathbf{b}} \cdot (I - X_0 X^{-1}) X \hat{\nabla} \hat{w} d\xi d\eta \right| \\ &= \left| \int_K v \mathbf{b} \cdot (I - X_0 X^{-1}) \nabla w dx dy \right| \\ &\leq Ch_K^\alpha \|\nabla w\|_{0,K} \|v\|_{0,K}, \end{aligned}$$

by Lemma 2.3.

$$(3.13) \quad \begin{aligned} & \left| \int_{\hat{K}} \hat{v} (\hat{\mathbf{b}} - \mathbf{b}_0) \cdot X_0 \hat{\nabla} \hat{w} d\xi d\eta \right| \\ &= \left| \int_{\hat{K}} \hat{v} (\hat{\mathbf{b}} - \mathbf{b}_0) \cdot X_0 X^{-1} (X \hat{\nabla} \hat{w}) d\xi d\eta \right| \\ &= \left| \int_K v (\mathbf{b} - \mathbf{b}_0) \cdot X_0 X^{-1} \nabla w dx dy \right| \\ &\leq Ch_K^\alpha \|\nabla w\|_{0,K} \|v\|_{0,K}, \end{aligned}$$

by Lemma 2.3 and (3.2). Applying (3.12) and (3.13) to (3.11), we obtain (3.6). \square

We then define two modified bilinear forms

$$a_h(u, v) = \sum_K a_h(u, v)_K, \quad b_h(w, v) = \sum_K b_h(w, v)_K$$

where

$$(3.14) \quad a_h(u, v)_K = (\nabla u, A \nabla v)_K^* + (\mathbf{b} \cdot \nabla u, v)_K + (cu, v)_K,$$

$$(3.15) \quad b_h(w, v)_K = (\nabla w, A \nabla v)_K^* + (\mathbf{b} \cdot \nabla w, v)_K + (cu, v)_K.$$

Given a quadrilateral partition \mathcal{T}_h on a polygonal domain Ω , we define the bilinear finite element space

$$S_h = \{v \in H^1(\Omega) : \hat{v} = v \circ F_K \in Q_1(\hat{K}), \forall K \in \mathcal{T}_h\}.$$

Theorem 3.2. Let \mathcal{T}_h satisfy the condition (α) and $RDP(N, \Psi)$, and let $u_I \in S_h$ be the bilinear interpolation of $u \in H^3(\Omega) \cap H_0^1(\Omega)$. Then there exists a constant C independent of h and u , such that for any $v \in S_h$,

$$|a_h(u - u_I, v)| + |b_h(u - u_I, v)| \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega})\|v\|_{1,\Omega}.$$

Proof: For convenience, we set $w = u - u_I$. By (3.14) and (3.15), we can express

$$a_h(w, v) - b_h(w, v) = \sum_{K \in \mathcal{T}_h} [(\mathbf{b} \cdot \nabla w, v)_K - (\mathbf{b} \cdot \nabla w, v)_K^*].$$

Recall (3.6), and we have

$$(3.16) \quad |a_h(w, v) - b_h(w, v)| \leq C \sum_{K \in \mathcal{T}_h} h_K^\alpha \|\nabla w\|_{0,K} \|v\|_{0,K} \\ \leq Ch^{1+\alpha} |u|_{2,\Omega} \|v\|_{0,\Omega},$$

since by the $RDP(N, \Psi)$ assumption, (2.14) is valid. Therefore, we only need to estimate $b_h(w, v)$. Again, by the $RDP(N, \Psi)$ assumption, we have

$$(3.17) \quad |(cw, v)| \leq Ch^2 |u|_{2,\Omega} \|v\|_{0,\Omega}.$$

Hence, our task is narrowed down to estimate

$$(\nabla u, A\nabla v)_K^*, \quad \text{and} \quad (\mathbf{b} \cdot \nabla w, v)_K^*$$

for $K \in \mathcal{T}_h$. By the definition (3.3) and (3.4), we see that all coefficients are constants now and we only need to estimate following terms

$$\int_{\hat{K}} \partial_\xi \hat{w} \partial_\xi \hat{v}, \quad \int_{\hat{K}} \partial_\xi \hat{w} \partial_\eta \hat{v}, \quad \int_{\hat{K}} \partial_\eta \hat{w} \partial_\xi \hat{v}, \quad \int_{\hat{K}} \partial_\eta \hat{w} \partial_\eta \hat{v}, \quad \int_{\hat{K}} \hat{v} \partial_\xi \hat{w}, \quad \int_{\hat{K}} \hat{v} \partial_\eta \hat{w}.$$

a) Let $\hat{u} \in P_2(\hat{K})$. There are only two terms ξ^2, η^2 not in the reference space of the bilinear interpolation, therefore,

$$\int_{\hat{K}} \partial_\xi \hat{w} \partial_\xi \hat{v} = 0, \quad \forall v \in S_h.$$

By the Bramble-Hilbert Lemma,

$$(3.18) \quad \left| \int_{\hat{K}} \partial_\xi \hat{w} \partial_\xi \hat{v} \right| \leq C \|D^3 \hat{u}\|_{L^2(\hat{K})} \|\partial_\xi \hat{v}\|_{L^2(\hat{K})} \\ \leq C (h_K^{1+\alpha} |u|_{2,K} + h_K^2 |u|_{3,K}) |v|_{1,K}.$$

We have used (2.2) and (2.3) in the last step. Similarly,

$$(3.19) \quad \left| \int_{\hat{K}} \partial_\eta \hat{w} \partial_\eta \hat{v} \right| \leq C (h_K^{1+\alpha} |u|_{2,K} + h_K^2 |u|_{3,K}) |v|_{1,K}.$$

Next we discuss the cross terms. For any $v \in S_h$, we can express

$$\partial_\xi \hat{v} = \partial_\xi \hat{v}(0, 0) + \eta \partial_{\xi\eta}^2 \hat{v}, \quad \partial_\eta \hat{v} = \partial_\eta \hat{v}(0, 0) + \xi \partial_{\xi\eta}^2 \hat{v}.$$

Note that $\partial_{\xi\eta}^2 \hat{v}$ is a constant. We write

$$\int_{\hat{K}} (\partial_\xi \hat{w} \partial_\eta \hat{v} \pm \partial_\eta \hat{w} \partial_\xi \hat{v}) \\ = \partial_\eta \hat{v}(0, 0) \int_{\hat{K}} \partial_\xi \hat{w} \pm \partial_\xi \hat{v}(0, 0) \int_{\hat{K}} \partial_\eta \hat{w} + \partial_{\xi\eta}^2 \hat{v} \left(\int_{\hat{K}} \xi \partial_\xi \hat{w} \pm \int_{\hat{K}} \eta \partial_\eta \hat{w} \right).$$

Since for $\hat{u} = \xi^2$, or $\hat{u} = \eta^2$,

$$\int_{\hat{K}} \partial_\xi \hat{w} = 0, \quad \int_{\hat{K}} \partial_\eta \hat{w} = 0.$$

Therefore, by the Bramble-Hilbert Lemma,

$$(3.20) \quad |\partial_\eta \hat{v}(0, 0) \int_{\hat{K}} \partial_\xi \hat{w} \pm \partial_\xi \hat{v}(0, 0) \int_{\hat{K}} \partial_\eta \hat{w}| \\ \leq C \|D^3 \hat{u}\|_{L^2(\hat{K})} \|\hat{\nabla} \hat{v}\|_{L^2(\hat{K})} \leq C (h_K^{1+\alpha} |u|_{2,K} + h_K^2 |u|_{3,K}) |v|_{1,K}.$$

Next we consider,

$$\begin{aligned}
\int_{\hat{K}} \xi \partial_\xi \hat{w} &= \frac{1}{2} \int_{\hat{K}} (\xi^2 - 1)' \partial_\xi \hat{w} = -\frac{1}{2} \int_{\hat{K}} (\xi^2 - 1) \partial_{\xi^2}^2 \hat{u}, \\
\partial_{\xi\eta}^2 \hat{v} \int_{\hat{K}} \xi \partial_\xi \hat{w} &= -\frac{1}{2} \int_{\hat{K}} (\xi^2 - 1) \partial_{\xi^2}^2 \hat{u} \partial_{\xi\eta}^2 \hat{v} \\
&= \frac{1}{2} \int_{-1}^1 (\xi^2 - 1) (\partial_{\xi^2}^2 \hat{u} \partial_\xi \hat{v})(\xi, -1) d\xi \\
&\quad - \frac{1}{2} \int_{-1}^1 (\xi^2 - 1) (\partial_{\xi^2}^2 \hat{u} \partial_\xi \hat{v})(\xi, 1) d\xi + \frac{1}{2} \int_{\hat{K}} (\xi^2 - 1) \partial_{\xi^2\eta}^3 \hat{u} \partial_\xi \hat{v}.
\end{aligned}$$

Similarly,

$$\begin{aligned}
\partial_{\xi\eta}^2 \hat{v} \int_{\hat{K}} \eta \partial_\eta \hat{w} &= \frac{1}{2} \int_{-1}^1 (\eta^2 - 1) (\partial_{\eta^2}^2 \hat{u} \partial_\eta \hat{v})(-1, \eta) d\eta \\
&\quad - \frac{1}{2} \int_{-1}^1 (\eta^2 - 1) (\partial_{\eta^2}^2 \hat{u} \partial_\eta \hat{v})(1, \eta) d\eta + \frac{1}{2} \int_{\hat{K}} (\eta^2 - 1) \partial_{\xi\eta^2}^3 \hat{u} \partial_\eta \hat{v}.
\end{aligned}$$

Therefore, we have

$$\begin{aligned}
(3.21) \quad \partial_{\xi\eta}^2 \hat{v} \left(\int_{\hat{K}} \xi \partial_\xi \hat{w} \pm \int_{\hat{K}} \eta \partial_\eta \hat{w} \right) &= \frac{1}{2} \int_{\partial\hat{K}}' (t^2 - 1) \partial_t^2 \hat{u} \partial_t \hat{v} dt \\
&\quad + \frac{1}{2} \int_{\hat{K}} [(\xi^2 - 1) \partial_{\xi^2\eta}^3 \hat{u} \partial_\xi \hat{v} \pm (\eta^2 - 1) \partial_{\xi\eta^2}^3 \hat{u} \partial_\eta \hat{v}],
\end{aligned}$$

where \int' indicates a sign influence whenever it applies. For the second term on the right hand side of (3.21), we have, from (2.3),

$$\begin{aligned}
(3.22) \quad &\frac{1}{2} \int_{\hat{K}} [(\xi^2 - 1) \partial_{\xi^2\eta}^3 \hat{u} \partial_\xi \hat{v} \pm (\eta^2 - 1) \partial_{\xi\eta^2}^3 \hat{u} \partial_\eta \hat{v}] \\
&\leq C(h_K^{1+\alpha} |u|_{2,K} + h_K^2 |u|_{3,K}) |v|_{1,K}.
\end{aligned}$$

In light of (3.18)–(3.22), we can express

$$\begin{aligned}
(3.23) \quad \int_{\hat{K}} (\hat{\nabla} \hat{w})^T B^K \hat{\nabla} \hat{v} d\xi d\eta &= \frac{b_{12}^K}{2} \sum_{j=1}^4 |l_j|^2 \int_{l_j} (t(s)^2 - 1) \partial_s^2 u \partial_s v ds \\
&\quad + (O(h_K^{1+\alpha}) |u|_{2,K} + O(h_K^2) |u|_{3,K}) |v|_{1,K},
\end{aligned}$$

where l_j are four sides of K . By the neighboring condition (2.13), any two adjacent elements K_1, K_2 that share a common edge satisfy (see Lemma 3.1)

$$\|B^{K_1} - B^{K_2}\| = O(h^\alpha),$$

Therefore, we have, by the trace theory,

$$\begin{aligned}
(3.24) \quad &\left| \frac{b_{12}^{K_1} - b_{12}^{K_2}}{2} |l|^2 \int_l (t(s)^2 - 1) \partial_s^2 u \partial_s v ds \right| \\
&\leq Ch^\alpha |l|^2 (h^{-1} \int_K |D^2 u Dv| + \int_K |D^3 u Dv + D^2 u D^2 v|) \\
&\leq C(h^{1+\alpha} |u|_{2,K} + h^2 |u|_{3,K}) |v|_{1,K}.
\end{aligned}$$

In the last step, we have used the inverse inequality. Adding up (3.23) with the edge integral estimated by (3.24), we obtain, under the homogeneous Dirichlet boundary condition,

$$(3.25) \quad \left| \sum_{K \in \mathcal{T}_h} (\nabla w, A \nabla v)_K^* \right| \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega})|v|_{1,\Omega}.$$

b) we now consider $\int_{\hat{K}} \hat{v} \partial_\xi \hat{w}$ where we can express

$$\hat{v} = \hat{v}(0,0) + \partial_\xi \hat{v}(0,0)\xi + \partial_\eta \hat{v}(0,0)\eta + \partial_{\xi\eta}^2 \hat{v} \xi \eta.$$

Since for any $\hat{u} \in P_2(\hat{K})$, we have

$$\int_{\hat{K}} \partial_\xi \hat{w} (\hat{v}(0,0) + \partial_\eta \hat{v}(0,0)\eta) = 0,$$

by the same argument as in b), we have

$$(3.26) \quad \begin{aligned} \left| \int_{\hat{K}} \partial_\xi \hat{w} (\hat{v}(0,0) + \partial_\eta \hat{v}(0,0)\eta) \right| &\leq C \|D^3 \hat{u}\|_{0,\hat{K}} \|\hat{v}\|_{1,\hat{K}} \\ &\leq C(h_K^\alpha |u|_{2,K} + h_K |u|_{3,K}) \|v\|_{1,K}. \end{aligned}$$

Next, by identities

$$\begin{aligned} \int_{\hat{K}} \partial_\xi \hat{w} \xi &= -\frac{1}{2} \int_{\hat{K}} \partial_{\xi^2}^2 \hat{u} (\xi^2 - 1), \\ \int_{\hat{K}} \partial_\xi \hat{w} \xi \eta &= -\frac{1}{4} \int_{\hat{K}} \partial_\xi \hat{w} (\xi^2 - 1)' (\eta^2 - 1)' = \frac{1}{4} \int_{\hat{K}} \partial_{\xi^2 \eta}^3 \hat{u} (\xi^2 - 1) (\eta^2 - 1), \end{aligned}$$

we have

$$(3.27) \quad \begin{aligned} \left| \int_{\hat{K}} \partial_\xi \hat{w} (\partial_\xi \hat{v}(0,0)\xi + \partial_{\xi\eta}^2 \hat{v} \xi \eta) \right| \\ \leq \|\partial_{\xi^2}^2 \hat{w}\|_{0,\hat{K}} \|\partial_\xi \hat{v}\|_{0,\hat{K}} + \|\partial_{\xi^2 \eta}^3 \hat{w}\|_{0,\hat{K}} \|\partial_{\xi\eta}^2 \hat{v}\|_{0,\hat{K}} \\ \leq C(h_K^\alpha |u|_{2,K} + h_K |u|_{3,K}) |v|_{1,K}. \end{aligned}$$

Again, we have used (2.2), (2.3), and the inverse inequality in the last step.

Combining (3.26) and (3.27), we obtain

$$(3.28) \quad \left| \int_{\hat{K}} \hat{v} \partial_\xi \hat{w} \right| \leq C(h_K^\alpha |u|_{2,K} + h_K |u|_{3,K}) \|v\|_{1,K}.$$

Similarly, we have

$$(3.29) \quad \left| \int_{\hat{K}} \hat{v} \partial_\eta \hat{w} \right| \leq C(h_K^\alpha |u|_{2,K} + h_K |u|_{3,K}) \|v\|_{1,K}.$$

Note that $X_0^K = O(h_K)$, therefore,

$$(3.30) \quad \begin{aligned} |(\mathbf{b}_0 \cdot \nabla w, v)_K^*| &= |\mathbf{b}_0 \cdot X_0 \int_{\hat{K}} \hat{v} \nabla \hat{w}| \\ &\leq Ch_K (h_K^\alpha |u|_{2,K} + h_K |u|_{3,K}) \|v\|_{1,K}. \end{aligned}$$

Adding up all $K \in \mathcal{T}_h$ and using the Cauchy inequality, we obtain

$$(3.31) \quad \left| \sum_{K \in \mathcal{T}_h} (\mathbf{b}_0 \cdot \nabla w, v)_K^* \right| \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega}) \|v\|_{1,\Omega}.$$

Combining (3.17), (3.25), and (3.31), we establish the assertion for $b_h(w, v)$.
□

Theorem 3.3. Assume that \mathcal{T}_h satisfies the condition (α) and $RDP(N, \Psi)$. Let $u \in H^3(\Omega) \cap H_0^1(\Omega)$ solves (3.1), let $u_h, u_I \in S_h$ be the finite element approximation and the bilinear interpolation of u , respectively, and let $a(\cdot, \cdot)$ satisfy the discrete inf-sup condition on S_h . Then there exists a constant C independent of h and u , such that

$$(3.32) \quad |a(u - u_I, v)| \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega})\|v\|_{1,\Omega},$$

$$(3.33) \quad \|u_h - u_I\|_{1,\Omega} \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega}).$$

Proof: Let $w = u - u_I$, and by Theorem 3.1,

$$\begin{aligned} |a(w, v)_K - a_h(w, v)_K| &= |(\nabla w, A\nabla v)_K - (\nabla w, A\nabla v)_K^*| \\ &\leq Ch_K^\alpha \|\nabla w\|_{0,K} \|\nabla v\|_{0,K}. \end{aligned}$$

Adding all $K \in \mathcal{T}_h$ and using (2.14) with the Cauchy-Schwarz inequality, we have

$$|a(w, v) - a_h(w, v)| \leq Ch^{1+\alpha}|u|_{2,\Omega}\|v\|_{1,\Omega}.$$

Recall Theorem 3.2, and we obtain

$$|a(w, v)| \leq |a(w, v) - a_h(w, v)| + |a_h(w, v)| \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega})\|v\|_{1,\Omega},$$

which establishes (3.32). We then complete the proof by the inf-sup condition in

$$\begin{aligned} c\|u_h - u_I\|_{1,\Omega} &\leq \sup_{v \in S_h} \frac{a(u_h - u_I, v)}{\|v\|_{1,\Omega}} \\ &= \sup_{v \in S_h} \frac{a(u - u_I, v)}{\|v\|_{1,\Omega}} \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega}). \quad \square \end{aligned}$$

4. Gradient Recovery

In this section, we introduce and analyze a polynomial preserving recovery method (PPR). We define a gradient recovery operator $G_h : S_h \rightarrow S_h \times S_h$, on bilinear finite element space under a quadrilateral partition \mathcal{T}_h in a following way: Given a finite element solution u_h , we first define $G_h u_h$ at all nodes (vertices), and then obtain $G_h u_h$ on the whole domain by interpolation using the original nodal shape functions of S_h .

Given an interior node (vertex) \mathbf{z}_i , we select an element patch ω_i , where

$$\bar{\omega}_i = \bigcup_{K \in \mathcal{T}_h, \mathbf{z}_i \in \bar{K}} \bar{K}.$$

We then denote all nodes on $\bar{\omega}_i$ (including \mathbf{z}_i) as \mathbf{z}_{ij} , $j = 1, 2, \dots, n(\geq 6)$, and fit a quadratic polynomial, in the least-squares sense, to the finite element solution u_h at those nodes. Using local coordinates (x, y) with \mathbf{z}_i as the origin, the fitting polynomial is

$$p_2(x, y; \mathbf{z}_i) = \mathbf{P}^T \mathbf{a} = \hat{\mathbf{P}}^T \hat{\mathbf{a}},$$

with

$$\mathbf{P}^T = (1, x, y, x^2, xy, y^2), \quad \hat{\mathbf{P}}^T = (1, \xi, \eta, \xi^2, \xi\eta, \eta^2);$$

$$\mathbf{a}^T = (a_1, a_2, a_3, a_4, a_5, a_6), \quad \hat{\mathbf{a}}^T = (a_1, ha_2, ha_3, h^2a_4, h^2a_5, h^2a_6),$$

where the scaling parameter $h = h_i$ is the length of the longest element edge in the patch ω_i . The coefficient vector $\hat{\mathbf{a}}$ is determined by the linear system

$$(4.1) \quad Q^T Q \hat{\mathbf{a}} = Q^T \mathbf{b}_h,$$

where $\mathbf{b}_h^T = (u_h(\mathbf{z}_{i1}), u_h(\mathbf{z}_{i2}), \dots, u_h(\mathbf{z}_{in}))$ and

$$Q = \begin{pmatrix} 1 & \xi_1 & \eta_1 & \xi_1^2 & \xi_1\eta_1 & \eta_1^2 \\ 1 & \xi_2 & \eta_2 & \xi_2^2 & \xi_2\eta_2 & \eta_2^2 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & \xi_n & \eta_n & \xi_n^2 & \xi_n\eta_n & \eta_n^2 \end{pmatrix}.$$

The condition for (4.1) to have a unique solution is: Q has a full rank, which is always satisfied in practical situations. In fact, Q has a full rank if and only if \mathbf{z}_{ij} s are not all lying on a conic curve. In practice, this is not a restriction at all: Any interior node \mathbf{z}_i is a common vertex of at least three quadrilaterals. This makes $n \geq 7 > 6$. As long as these n nodes are reasonably distributed, it is impossible to fit all of them on a conic curve. A simple example is that three nodes are on the same line and others are not form a line.

Conjecture: A sufficient condition for Q to have a full rank is all quadrilaterals are convex.

In the rest of this article, we will always assume the *rank condition*: Q has a full rank.

Now we define

$$(4.2) \quad G_h u_h(\mathbf{z}_i) = \nabla p_2(0, 0; \mathbf{z}_i).$$

When Neumann boundary condition is post, there is no need to do gradient recovery on the boundary. However, if the Dirichlet boundary condition is post, the recovered gradient on a boundary node \mathbf{z} can be determined from an element patch ω_i such that $\mathbf{z} \in \bar{\omega}_i$ in the following way: Let the relative coordinates of \mathbf{z} with respect to \mathbf{z}_i is, say (h, h) , then $G_h u_h(\mathbf{z}) = \nabla p_2(h, h; \mathbf{z}_i)$. If \mathbf{z} is covered by more than one element patches, then some averaging may be applied.

Remark 4.1. In an earlier work [26], Wiberg-Li least-squares fitted solution values to improve and to estimate the L_2 -norm errors of the finite element approximation.

Now, we demonstrate PPR on an element patch that contains four uniform square elements (Figure 2). Fitting

$$\hat{p}_2(\xi, \eta) = (1, \xi, \eta, \xi^2, \xi\eta, \eta^2)(\hat{a}_1, \dots, \hat{a}_6)^T$$

with respect to the nine nodal values on the patch. Now

$$\vec{e} = (1, 1, 1, 1, 1, 1, 1, 1, 1)^T, \quad \vec{\xi} = (0, 1, 1, 0, -1, -1, -1, 0, 1)^T,$$

$$\vec{\eta} = (0, 0, 1, 1, 1, 0, -1, -1, -1)^T, \quad Q = (\vec{e}, \vec{\xi}, \vec{\eta}, \vec{\xi}^2, \vec{\xi}\vec{\eta}, \vec{\eta}^2),$$




Figure 2

$$(Q^T Q)^{-1} Q^T = \text{diag}\left(\frac{1}{9}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{4}, \frac{1}{6}\right) \cdot \begin{pmatrix} 5 & 2 & -1 & 2 & -1 & 2 & -1 & 2 & -1 \\ 0 & 1 & 1 & 0 & -1 & -1 & -1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & 0 & -1 & -1 & -1 \\ -2 & 1 & 1 & -2 & 1 & 1 & 1 & -2 & 1 \\ 0 & 0 & 1 & 0 & -1 & 0 & 1 & 0 & -1 \\ -2 & -2 & 1 & 1 & 1 & -2 & 1 & 1 & 1 \end{pmatrix}.$$

(4.3)

$$G_h u(\mathbf{z}) = \frac{1}{h} \hat{\nabla} \hat{p}_2(0, 0) = \frac{1}{6h} \begin{pmatrix} u_1 - u_5 + u_2 - u_4 + u_8 - u_6 \\ u_2 - u_8 + u_3 - u_7 + u_4 - u_6 \end{pmatrix} = \frac{1}{h} \sum_j \vec{c}_j u_j.$$

Note that the desired weights \vec{c}_j are the second row of $(Q^T Q)^{-1} Q^T$ for the x -derivative, and the third row of $(Q^T Q)^{-1} Q^T$ for the y -derivative, respectively. Moreover, $\sum_i \vec{c}_j = \vec{0}$ and $G_h u(\mathbf{z})$ provides a second-order finite

difference scheme at \mathbf{z} .

Given $v \in S_h$, it is straightforward to verify that

$$\begin{aligned} \frac{\partial v}{\partial x}\left(\frac{h}{2}, \frac{2h}{3}\right) &= \frac{1}{3h}(v_1 - v_0) + \frac{2}{3h}(v_2 - v_3), \\ \frac{\partial v}{\partial x}\left(-\frac{h}{2}, \frac{2h}{3}\right) &= \frac{1}{3h}(v_0 - v_5) + \frac{2}{3h}(v_3 - v_4), \\ \frac{\partial v}{\partial x}\left(-\frac{h}{2}, -\frac{2h}{3}\right) &= \frac{1}{3h}(v_0 - v_5) + \frac{2}{3h}(v_7 - v_6), \end{aligned}$$

$$\frac{\partial v}{\partial x}\left(\frac{h}{2}, -\frac{2h}{3}\right) = \frac{1}{3h}(v_1 - v_0) + \frac{2}{3h}(v_8 - v_7).$$

Therefore,

$$G_h^x v(\mathbf{z}) = \frac{1}{4}\left[\frac{\partial v}{\partial x}\left(\frac{h}{2}, \frac{2h}{3}\right) + \frac{\partial v}{\partial x}\left(-\frac{h}{2}, \frac{2h}{3}\right) + \frac{\partial v}{\partial x}\left(-\frac{h}{2}, -\frac{2h}{3}\right) + \frac{\partial v}{\partial x}\left(\frac{h}{2}, -\frac{2h}{3}\right)\right].$$

The recovered y -derivative can be obtained similarly. Hence, in this special case,

$$|G_h v(\mathbf{z})| \leq |v|_{1,\infty,\omega_{\mathbf{z}}}.$$

By linear mapping, this is also valid for four uniform parallelograms in that

$$(4.4) \quad |G_h v(\mathbf{z})| \leq C|v|_{1,\infty,\omega_{\mathbf{z}}}, \quad \forall v \in S_h.$$

with C independent of h and v .

Theorem 4.1 Let \mathcal{T}_h satisfy *Condition* (α). Then the recovery operator G_h is a bounded linear operator on bilinear element space such that

$$\|G_h v\|_{0,p,\Omega} \leq C|v|_{1,p,\Omega}, \quad \forall v \in S_h, \quad 1 \leq p \leq \infty,$$

where C is a constant independent of v and h .

Proof: We observe that the diagonal condition together with the neighboring condition imply that for any given node \mathbf{z} , there are four elements attached to it when h is sufficiently small. In addition, these four elements deviate from four parallelograms that attached to the same node in the following sense,

$$Q = Q_0 + h^\alpha Q_1,$$

where Q and Q_0 are least-square fitting matrices associated with those four quadrilateral elements and four parallelograms, respectively. We want to express $(Q^T Q)^{-1} Q^T$ in terms of $(Q_0^T Q_0)^{-1} Q_0^T$. Towards this end, we have

$$Q^T Q = Q_0^T Q_0 (I + h^\alpha E_1),$$

where

$$E_1 = (Q_0^T Q_0)^{-1} (Q_1^T Q_0 + Q_0^T Q_1 + h^\alpha Q_1^T Q_1).$$

Therefore,

$$(Q^T Q)^{-1} Q^T = (I + h^\alpha E_1)^{-1} (Q_0^T Q_0)^{-1} (Q_0^T + h^\alpha Q_1^T) = (Q_0^T Q_0)^{-1} Q_0^T + h^\alpha E_2,$$

where

$$E_2 = (Q_0^T Q_0)^{-1} Q_1^T - \sum_{j=0}^{\infty} (h^\alpha E_1)^j E_1 (Q_0^T Q_0)^{-1} Q_0^T.$$

We see that

$$(4.5) \quad (Q^T Q)^{-1} Q^T = (Q_0^T Q_0)^{-1} Q_0^T + O(h^\alpha).$$

Therefore, the fact that $Q_0^T Q_0$ is invertible guarantees that $Q^T Q$ is invertible for sufficiently small h . Moreover, by (4.5), we have

$$G_h v(\mathbf{z}) = \frac{1}{h} \sum_j (\vec{c}_j + O(h^\alpha)) v_j$$

where G_h is the recovery operator under the quadrilateral mesh that satisfies the *diagonal condition* and the *neighboring condition*, and \vec{c}_j s are weights

for the related parallelogram mesh so that, by (4.4), $\frac{1}{h} \sum_j \vec{c}_j$ is a bounded operator on S_h such that

$$\left| \frac{1}{h} \sum_j \vec{c}_j v_j \right| \leq C |v|_{1,\infty,\omega_{\mathbf{z}}}.$$

Therefore, in the quadrilateral case, (4.4) is also valid, provided *Condition* (α) is satisfied and h is sufficiently small. If (4.4) is valid for each node of K , then we have,

$$(4.6) \quad \|G_h v\|_{0,\infty,K} \leq C |v|_{1,\infty,\omega_K}, \quad \forall v \in S_h,$$

where ω_K is defined as

$$\bar{\omega}_K = \bigcup_{K' \in \mathcal{T}_h, \bar{K}' \cap \bar{K} \neq \emptyset} \bar{K}'.$$

Note that (4.6) is true for all $K \in \mathcal{T}_h$ including boundary elements, since by our construction the boundary recovery is simply some averaging of nearby patches. Therefore,

$$(4.7) \quad \|G_h v\|_{0,\infty,\Omega} \leq C |v|_{1,\infty,\Omega}, \quad \forall v \in S_h.$$

This establishes the assertion for $p = \infty$. As for $p < \infty$, we notice that all norms are equivalent for finite dimensional spaces, and with a scaling argument,

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_K |G_h v|^p &\leq C_1 \sum_{K \in \mathcal{T}_h} h^2 \|G_h v\|_{0,\infty,K}^p \\ &\leq C_2 h^2 \sum_{K \in \mathcal{T}_h} |v|_{1,\infty,K}^p \\ &\leq C_3 h^2 \sum_{K \in \mathcal{T}_h} h^{-2} \int_K |\nabla v|^p \leq C \sum_{K \in \mathcal{T}_h} |v|_{1,p,K}^p \end{aligned}$$

Here, all constants C_j 's and C are independent of p , v , and h . The conclusion then follows. \square

Another important feature of the new recovery operator is the following polynomial preserving property:

Lemma 4.1. Let $K \in \mathcal{T}_h$ and u be a quadratic polynomial on ω_K . Assume that Q has a full rank. Then $G_h u = \nabla u$ on K .

Proof: The rank condition guarantees that the least-squares fitting has a unique solution. On each of four element patches, the recovery procedure results in a quadratic polynomial p_2 that least-squares fits u , a quadratic polynomial. Therefore, $p_2 = u$, and consequently, $G_h u = \nabla p_2 = \nabla u$, a linear function, at each of the four vertices of K . Therefore, $G_h u = \nabla u$ on K . \square

Remark 4.2. Note that we do not make any mesh assumptions in Lemma 4.1 except the rank condition, which is always satisfied in practice. Basically,

as long as the least-squares fitting procedure can be carried out, the polynomial preserving property is satisfied. As a comparison, the ZZ recovery operator does not have this polynomial preserving property under general meshes, see [33] for more details.

Theorem 4.2. Let \mathcal{T}_h satisfy the condition (α) and $RDP(N, \Psi)$. Let $u_h \in S_h$ be the finite element approximation of $u \in H^3(\Omega) \cap H_0^1(\Omega)$, the solution of (3.1), and let $a(\cdot, \cdot)$ satisfy the discrete inf-sup condition on S_h . Then the recovered gradient is superconvergent in the sense

$$\|\nabla u - G_h u_h\|_{0,\Omega} \leq C(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega}),$$

where C is a constant independent of u and h .

Proof: We decompose the error into

$$(4.8) \quad \nabla u - G_h u_h = \nabla u - G_h u + G_h(u_I - u_h).$$

Note that $G_h u = G_h u_I$ since $u_I = u$ at all vertices and the recovery operator G_h is completely determined by nodal values of u . By the polynomial preserving property and the Bramble-Hilbert lemma,

$$(4.9) \quad \|\nabla u - G_h u\|_{0,p,\Omega} \leq Ch^2|u|_{3,p,\Omega}, \quad 1 \leq p \leq \infty.$$

By Theorem 4.1, G_h is a bounded operator for all interior patches. Therefore,

$$(4.10) \quad \begin{aligned} \|G_h(u_I - u_h)\|_{0,\Omega}^2 &= \sum_{K \in \mathcal{T}_h} \|G_h(u_I - u_h)\|_{0,K}^2 \\ &\leq C^2 \sum_{K \in \mathcal{T}_h} |u_I - u_h|_{1,K}^2 \leq C^2(h^{1+\alpha}|u|_{2,\Omega} + h^2|u|_{3,\Omega})^2 \end{aligned}$$

by Theorem 3.3. The conclusion then follows by applying (4.9) with $p = 2$ and (4.10) to (4.8). \square

Theorem 4.2 assumes a global regularity $u \in H^3(\Omega)$, which may not hold in general. However, higher regularity requirement is usually satisfied in an interior sub-domain. In the rest of this section, we shall prove a local result based on interior estimates. In order to concentrate on superconvergence analysis, treatments of curved boundaries and corner singularities will not be discussed here. We merely assume that they have been taken care of in the following sense,

$$(4.11) \quad \|u - u_h\|_{-1,\Omega} \leq C(f, a, \Omega)h^{1+\rho}, \quad \rho = \min(1, \alpha).$$

The negative norm term is the only one in our analysis that takes into account what happens outside of a local region Ω_1 .

We shall show that under assumption (4.11), superconvergent recovery will occur in an interior sub-domain. Toward this end, we consider $\Omega_0 \subset\subset \Omega_1 \subset\subset \Omega$ where Ω_0 and Ω_1 are compact polygonal sub-domains that can be decomposed into quadrilaterals. By ‘‘compact sub-domains’’ we mean that $dist(\Omega_0, \partial\Omega_1)$ and $dist(\Omega_1, \partial\Omega)$ are of order $O(1)$. Outside Ω_1 , we may have quadrilateral or triangular subdivisions. We may also have refined meshes near the corner singularities and curved elements on the boundary regions. We assume that all these together will result in (4.11).

We define a cut-off function $\omega \in C_0^\infty(\Omega)$ such that $\omega = 1$ on Ω_0 and $\omega = 0$ in $\Omega \setminus \Omega_1$. We decompose u into

$$u = \tilde{u} + \hat{u}, \quad \tilde{u} = u\omega.$$

Let \tilde{u}_I be the bilinear interpolation of \tilde{u} and let $\tilde{u}_h \in S_h(\Omega_1) = S_h \cap H_0^1(\Omega_1)$ be the finite element approximation of \tilde{u} on Ω_1 . There holds

$$a(\tilde{u} - \tilde{u}_h, v)_{\Omega_1} = 0, \quad \forall v \in S_h(\Omega_1).$$

The index Ω_1 indicates that the integrations in the bilinear form are performed on the subdomain. Further, we let

$$\hat{u}_I = u_I - \tilde{u}_I, \quad \hat{u}_h = u_h - \tilde{u}_h.$$

Note that we have $\tilde{u}_I = u_I$ on Ω_0 since $\omega = 1$ on Ω_0 . However, $\tilde{u}_h \neq u_h$ on Ω_0 in general.

Apply Theorem 3.3 on Ω_1 , and we immediately obtain

$$(4.12) \quad \|\tilde{u}_h - \tilde{u}_I\|_{1,\Omega_1} \leq Ch^{1+\rho} \|\tilde{u}\|_{3,\Omega_1}.$$

However, for any $k \leq 3$,

$$(4.13) \quad |\tilde{u}|_{k,\Omega_1} = |u\omega|_{k,\Omega_1} \leq \sum_{j=0}^k |D^j u D^{k-j} \omega|_{L_2(\Omega_1)} \leq C(k, \omega) \|u\|_{k,\Omega_1}.$$

Therefore, from (4.12),

$$(4.14) \quad \|\tilde{u}_h - \tilde{u}_I\|_{1,\Omega_0} \leq \|\tilde{u}_h - \tilde{u}_I\|_{1,\Omega_1} \leq Ch^{1+\rho} \|u\|_{3,\Omega_1}.$$

Next, we consider $\hat{u}_h - \hat{u}_I$. Since $\hat{u} = \hat{u}_I = 0$ on Ω_0 , there holds

$$(4.15) \quad \|\hat{u}_h - \hat{u}_I\|_{1,\Omega_0} = \|\hat{u}_h\|_{1,\Omega_0} = \|\hat{u}_h - \hat{u}\|_{1,\Omega_0}.$$

Note that for all $v \in S_h(\Omega_1)$,

$$a(\hat{u}_h - \hat{u}, v)_{\Omega_1} = a(u_h - u, v)_{\Omega_1} - a(\tilde{u}_h - \tilde{u}, v)_{\Omega_1} = 0.$$

As a result,

$$(4.16) \quad \|\hat{u} - \hat{u}_h\|_{1,\Omega_0} \leq C(h^2 \|\hat{u}\|_{3,\Omega_1} + \|\hat{u} - \hat{u}_h\|_{-1,\Omega_1}),$$

by Nitsche and Schatz [20, Theorem 5.1] (All the conditions of this theorem can be verified in the current situation, see Remark 4.3 below).

With the same argument as in (4.13), we have

$$(4.17) \quad \|\hat{u}\|_{3,\Omega_1} = \|u - \tilde{u}\|_{3,\Omega_1} \leq \|u\|_{3,\Omega_1} + \|\tilde{u}\|_{3,\Omega_1} \leq C \|u\|_{3,\Omega_1}.$$

Observe that

$$\|\tilde{u} - \tilde{u}_h\|_{-1,\Omega_1} \leq \|\tilde{u} - \tilde{u}_h\|_{0,\Omega_1} \leq Ch^2 \|u\|_{2,\Omega_1},$$

therefore, by assumption (4.11),

$$(4.18) \quad \begin{aligned} \|\hat{u} - \hat{u}_h\|_{-1,\Omega_1} &\leq \|u - u_h\|_{-1,\Omega_1} + \|\tilde{u} - \tilde{u}_h\|_{-1,\Omega_1} \\ &\leq Ch^{1+\rho} (C(f, a, \Omega) + \|u\|_{2,\Omega_1}). \end{aligned}$$

Substituting (4.17) and (4.18) into (4.16), we derive

$$(4.19) \quad \|\hat{u} - \hat{u}_h\|_{1,\Omega_0} \leq Ch^{1+\rho} (\|u\|_{3,\Omega_1} + C(f, a, \Omega)).$$

Combining (4.19) with (4.14) and (4.15), we obtain

$$(4.20) \quad \begin{aligned} \|u_h - u_I\|_{1,\Omega_0} &\leq \|\tilde{u}_h - \tilde{u}_I\|_{1,\Omega_0} + \|\hat{u}_h - \hat{u}_I\|_{1,\Omega_0} \\ &\leq Ch^{1+\rho}(\|u\|_{3,\Omega_1} + C(f, a, \Omega)). \end{aligned}$$

Now, following the same argument as in Theorem 4.2, we immediately obtain the following result on a general polygonal domain.

Theorem 4.3. Let $\Omega \subset R^2$ be a polygonal domain and $\Omega_0 \subset\subset \Omega_1 \subset\subset \Omega$. Assume that \mathcal{T}_h satisfy the condition (α) and $RDP(N, \Psi)$ on Ω_1 . Let $u_h \in S_h$ be the finite element approximation of $u \in H^3(\Omega_1) \cap H_0^1(\Omega)$ that solves (3.1) with $a(\cdot, \cdot)$ satisfying the discrete inf-sup condition on S_h . Furthermore, let (4.11) be satisfied. Then there exists a constant C independent of h such that

$$\|G_h u_h - \nabla u\|_{1,\Omega_0} \leq Ch^{1+\rho}(\|u\|_{3,\Omega_1} + C(f, a, \Omega)), \quad \rho = \min(1, \alpha).$$

Remark 4.3. In the proof of Theorem 4.3, we used a result of Nitsche and Schatz [20, Theorem 5.1], which requires following conditions from the underlining finite element space:

- R1. Coercive and continuity of the bilinear form;
- A.1. Approximation in an optimal sense;
- A.2. Superapproximation property;
- A.3. Inverse properties.

In our situation, R1 is assured by the inf-sup condition and A.1. is actually (2.14). Here we sketch a proof for A.2. under a quadrilateral mesh. The verification for A.3. is similar.

We want to show that for $\Omega_0 \subset\subset G \subset\subset \Omega_1$, $v_h \in S_h$ and $\omega \in C_0^\infty(\Omega_0)$, there exists an $\eta \in S_h(G)$ such that

$$(4.21) \quad \|\omega v_h - \eta\|_{1,G} \leq Ch\|v_h\|_{1,G}.$$

By direct calculation over a quadrilateral element K , we have

$$\begin{aligned} |\omega v_h - \eta|_{1,K}^2 &= \int_{\hat{K}} \frac{1}{J_K} |X \hat{\nabla}(\widehat{\omega v_h} - \hat{\eta})|^2 d\xi d\eta \\ &\leq C \left(\left\| \frac{\partial^2}{\partial \xi^2}(\widehat{\omega v_h}) \right\|_{L^2(\hat{K})}^2 + \left\| \frac{\partial^2}{\partial \eta^2}(\widehat{\omega v_h}) \right\|_{L^2(\hat{K})}^2 \right) \leq Ch_K^2 \|v_h\|_{H^1(K)}^2. \end{aligned}$$

Indeed, since $\partial_{\xi^2} \hat{v}_h = 0$ for a bilinear function on \hat{K} , we have

$$\begin{aligned} \frac{\partial^2}{\partial \xi^2}(\widehat{\omega v_h}) &= \frac{\partial^2 \hat{\omega}}{\partial \xi^2} v_h + 2 \frac{\partial \hat{\omega}}{\partial \xi} \frac{\partial \hat{v}_h}{\partial \xi} \\ &= v_h [(a_1 + a_3 \eta) \partial_x + (b_1 + b_3 \eta) \partial_y]^2 \omega + 2 DF_K \nabla \omega \cdot DF_K \nabla v_h. \end{aligned}$$

We then have the needed power for h_K . Another term $\|\omega v_h - \eta\|_{L^2(K)}$ can be similarly estimated. Finally, we simply add up $K \subset G$ and taking the square root to obtain (4.21).

5. A Posteriori Error Estimates

Let $e_h = u - u_h$, the task here is to estimate the error $\|\nabla e_h\|_{0,\Omega_0}$ by a computable quantity η_h . According to Zienkiewicz-Zhu [36], η_h is the error estimator defined by the recovered gradient,

$$\eta_h = \|G_h u_u - \nabla u_h\|_{0,\Omega_0}.$$

We need the following assumption:

$$(5.1) \quad \|\nabla e_h\|_{0,\Omega_0} \geq Ch.$$

Theorem 5.1. Assume the same hypotheses as in Theorem 4.3. Let (5.1) be satisfied. Then

$$\frac{\eta_h}{\|\nabla e_h\|_{0,\Omega_0}} = 1 + O(h^\rho), \quad \rho = \min(1, \alpha).$$

Proof: By the triangle inequality,

$$\eta_h - \|\nabla u - G_h u_h\|_{0,\Omega_0} \leq \|\nabla e_h\|_{0,\Omega_0} \leq \eta_h + \|\nabla u - G_h u_h\|_{0,\Omega_0}.$$

Dividing the above by $\|\nabla e_h\|_{0,\Omega_0}$, the conclusion follows from Theorem 4.3 and (5.1). \square

Remark 5.1. Theorem 5.1 indicates that the error estimator based on our polynomial preserving recovery is asymptotically exact on an interior region Ω_0 . This result is valid for fairly general quadrilateral meshes.

Remark 5.2. If we use $o(1)$ to substitute $O(h^\alpha)$, the conclusion of Theorem 5.1 would be $1 + o(1)$ and the error estimate would still be asymptotically correct. Furthermore, we may use a more practical term: “a sufficiently small constant $\gamma > 0$ ”, instead of $o(1)$. We would lose the asymptotic exactness, nevertheless, the effectivity index would still be in a reasonable range around 1, as observed in practice.

Appendix

Proof of Lemma 2.1. Let the longest edge length of K be h_K , then the longest edge length after one bisection refinement is $h_K/2$. We shall show that the distance between the two diagonal mid-points of any one of the four refined quadrilaterals is $d_K/4$, a quadratic reduction.

1) The coordinates O show that P_1P_3 , P_4P_2 and O_1O_2 bisect each other at O .

2) $|O_1P_2| = |Z_3Z_4|/2 = |Z_3P_3|$ since O_1P_2 connects two edge centers in $\Delta Z_2Z_3Z_4$.

3) $|Z_3Q_1| = |Q_1O_1|$ since two triangles $\Delta Q_1O_1P_2$ and $\Delta Q_1Z_3P_3$ are congruent.

4) $|Q_1Q_2| = |OO_1|/2 = d_K/4$ since Q_1Q_2 connects two edge centers in ΔZ_3OO_1 . \square

Proof of Lemma 2.2. From

$$\begin{aligned} |\mathbf{v}_1||\mathbf{v}_3|\sin(\pi - \theta_1) &= |\mathbf{v}_1 \times \mathbf{v}_3| = \frac{1}{2}|J_1^K| \\ &= \frac{1}{8}|P_4P_2 \times O_1O_2| = \frac{1}{8}|P_4P_2|d_K \sin \beta_K, \end{aligned}$$

we have

$$\sin(\pi - \theta_1) = \frac{|P_4 P_2 \times O_1 O_2|}{8|\mathbf{v}_1||\mathbf{v}_3|} = \frac{|P_4 P_2| d_K}{8|\mathbf{v}_1||\mathbf{v}_3|} \sin \beta_K.$$

Similarly,

$$\sin(\pi - \theta_2) = \frac{|P_1 P_3| d_K}{8|\mathbf{v}_2||\mathbf{v}_4|} \sin \gamma_K.$$

Note that

$$\begin{aligned} \min(|\mathbf{v}_1|, |\mathbf{v}_3|) &\leq |P_4 P_2| \leq \max(|\mathbf{v}_1|, |\mathbf{v}_3|), \\ \min(|\mathbf{v}_2|, |\mathbf{v}_4|) &\leq |P_1 P_3| \leq \max(|\mathbf{v}_2|, |\mathbf{v}_4|); \end{aligned}$$

and when σ_K is small,

$$\sin(\pi - \theta_1) \approx \pi - \theta_1, \quad \sin(\pi - \theta_2) \approx \pi - \theta_2.$$

The conclusion then follows. \square

References

- [1] G. Acosta and R.G. Durán, Error estimates for Q_1 isoparametric elements satisfying a weak angle condition, *SIAM J. Numer. Anal.* 38 (2001), 1073-1088.
- [2] M. Ainsworth and J.T. Oden, *A Posteriori Error Estimation in Finite Element Analysis*, Wiley Interscience, New York, 2000.
- [3] D.N. Arnold, D. Boffi, and R.S. Falk, Approximation by quadrilateral finite elements, *Math. Comp.* 71 (2002), 909-922.
- [4] I. Babuška and T. Strouboulis, *The Finite Element Method and its Reliability*, Oxford University Press, London, 2001.
- [5] R. Bank, Hierarchical bases and the finite element method, *Acta Numerica* (1996), 1-43.
- [6] R.E. Bank and J. Xu, Asymptotically exact *a posteriori* error estimators, Part I: Grids with superconvergence, *SIAM J. Numer. Anal.* 41 (2004), 2294-2312.
- [7] R.E. Bank and J. Xu, Asymptotically exact *a posteriori* error estimators, Part II: General unstructured grids, *SIAM J. Numer. Anal.* 41 (2004), 2313-2332.
- [8] S.C. Brenner and L.R. Scott, *The Mathematical Theory of Finite Element Methods*, Springer-Verlag, New York, 1994.
- [9] C. Carstensen, All first-order averaging techniques for a posteriori finite element error control on unstructured grids are efficient and reliable, to appear in *Math. Comp.*
- [10] C. Carstensen and S. Bartels, Each averaging technique yields reliable *a posteriori* error control in FEM on unstructured grids. Part I: Low order conforming, nonconforming, and mixed FEM, *Math. Comp.* 71 (2002), 945-969.
- [11] C. Chen and Y. Huang, *High Accuracy Theory of Finite Element Methods* (in Chinese), Hunan Science Press, P.R. China, 1995.
- [12] Y. Chen, Superconvergent recovery of gradients of piecewise linear finite element approximations on non-uniform mesh partitions, *Numer. Meths. PDEs* 14 (1998), 169-192.
- [13] P.G. Ciarlet, *The finite element method for elliptic problems*, North-Holland, Amsterdam, 1978.
- [14] V. Girault and P.-A. Raviart, *Finite element methods for Navier-Stokes equations*, Springer-Verlag, New York, 1986.
- [15] W. Hoffmann, A.H. Schatz, L.B. Wahlbin, and G. Wittum, Asymptotically exact *a posteriori* estimators for the pointwise gradient error on each element in irregular meshes. Part 1: A smooth problem and globally quasi-uniform meshes, *Math. Comp.* 70 (2001), 897-909.
- [16] M. Křížek, P. Neittaanmäki, and R. Stenberg (Eds.), *Finite Element Methods: Superconvergence, Post-processing, and A Posteriori Estimates*, Lecture Notes in Pure and Applied Mathematics Series, Vol. 196, Marcel Dekker, Inc., New York, 1997.

- [17] B. Li and Z. Zhang, Analysis of a class of superconvergence patch recovery techniques for linear and bilinear finite elements, *Numer. Meth. PDEs* 15 (1999), 151–167.
- [18] Q. Lin and N. Yan, *Construction and Analysis of High Efficient Finite Elements* (in Chinese), Hebei University Press, P.R. China, 1996.
- [19] P.B. Ming and Z.-C. Shi, Quadrilateral mesh revisited, *Comput. Methods Appl. Mech. Engrg.*, 191 (2002), 5671–5682.
- [20] J.A. Nitsche and A.H. Schatz, Interior estimates for Ritz-Galerkin methods, *Math. Comp.* 28 (1974), 937-958.
- [21] R. Rannacher and S. Turek, Simple nonconforming quadrilateral Stokes element, *Numer. Meth. PDEs* 8 (1992), 97-111.
- [22] A.H. Schatz and L.B. Wahlbin, Asymptotically exact *a posteriori* estimators for the pointwise gradient error on each element in irregular meshes. Part 2: The piecewise linear case, *Math. Comp.* 73 (2004), 517-523.
- [23] R. Verfürth, *A Review of A Posteriori Error Estimation and Adaptive Mesh-Refinement Techniques*, Wiley-Teubner, Stuttgart, 1996.
- [24] L.B. Wahlbin, Local behavior in finite element methods, in *Handbook of Numerical Analysis, Vol. II, Finite Element Methods* (Part 1), P.G. Ciarlet and J.L. Lions, ed., Elsevier Science Publishers B.V. (North-Holland), 1991, 353-522.
- [25] L.B. Wahlbin, *Superconvergence in Galerkin Finite Element Methods*, Lecture Notes in Mathematics, Vol. 1605, Springer, Berlin, 1995.
- [26] N.-E. Wiberg and X.D. Li, Superconvergence patch recovery of finite element solutions and a *a posteriori* L_2 norm error estimate, *Commun. Num. Meth. Eng.* 37 (1994), 313–320.
- [27] J. Wang, A superconvergence analysis for finite element solutions by the least-squares surface fitting on irregular meshes for smooth problems, *J. Math. Study* 33 (2000), 229-243.
- [28] J. Xu and Z. Zhang, Analysis of recovery type *a posteriori* error estimators for mildly structured grids, to appear in *Math. Comp.*
- [29] N. Yan and A. Zhou, Gradient recovery type *a posteriori* error estimates for finite element approximations on irregular meshes, *Comput. Methods Appl. Mech. Engrg.* 190 (2001), 4289-4299.
- [30] J. Zhang and F. Kikuchi, Interpolation error estimates of a modified 8-node serendipity finite element, *Numer. Math.* 85 (2000), 503-524.
- [31] Z. Zhang, Analysis of some quadrilateral nonconforming elements for incompressible elasticity, *SIAM J. Numer. Anal.* 34 (1997), 640-663.
- [32] Z. Zhang, Polynomial preserving recovery for anisotropic and irregular grids, *Journal of Computational Mathematics* 22 (2004), 331-340.
- [33] Z. Zhang and A. Naga, A new finite element gradient recovery method: Superconvergence property, Accepted for publication by *SIAM Journal on Scientific Computing*.
- [34] Q.D. Zhu and Q. Lin, *Superconvergence Theory of the Finite Element Method* (in Chinese), Hunan Science Press, China, 1989.
- [35] O.C. Zienkiewicz and J.Z. Zhu, The superconvergence patch recovery and *a posteriori* error estimates. Part 1: The recovery technique, *Int. J. Numer. Methods Engrg.* 33 (1992), 1331-1364.
- [36] O.C. Zienkiewicz and J.Z. Zhu, The superconvergent patch recovery and *a posteriori* error estimates, Part 2: Error estimates and adaptivity, *Int. J. Numer. Methods Engrg.* 33 (1992), 1365-1382.
- [37] M. Zlámal, Superconvergence and reduced integration in the finite element method, *Math. Comp.* 32 (1978), 663-685.

Department of Mathematics, Wayne State University, Detroit, MI 48202
E-mail: z Zhang@math.wayne.edu